

World Journal of Gastroenterology®

Volume 15 Number 19
May 21, 2009



Published by The WJG Press and Baishideng
Room 903, Building D, Ocean International Center,
No. 62 Dongsihuan Zhonglu, Chaoyang District, Beijing 100025, China
Telephone: +86-10-59080039
Fax: +86-10-85381893
E-mail: wjg@wjgnet.com
http://www.wjgnet.com

ISSN 1007-9327



World Journal of Gastroenterology

www.wjgnet.com

Volume 15

Number 19

May 21

2009



World Journal of Gastroenterology®

Indexed and Abstracted in:

Current Contents®/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch®) and Journal Citation Reports/Science Edition, *Index Medicus*, MEDLINE and PubMed, Chemical Abstracts, EMBASE/Excerpta Medica, Abstracts Journals, PubMed Central, Digital Object Identifier, CAB Abstracts and Global Health.
ISI JCR 2003-2000 IF: 3.318, 2.532, 1.445 and 0.993.

**Volume 15 Number 19
May 21, 2009**

World J Gastroenterol
2009 May 21; 15(19): 2305-2432

Online Submissions

wjg.wjgnet.com
www.wjgnet.com

Printed on Acid-free Paper

世界胃肠病学杂志



World Journal of Gastroenterology®

Weekly Established in October 1995

Volume 15 Number 19
May 21, 2009



Contents

EDITORIAL	2305	Hepatitis C virus lymphotropism and peculiar immunological phenotype: Effects on natural history and antiviral therapy <i>Conca P, Tarantino G</i>
	2309	Clinicopathological features of early gastric cancer with duodenal invasion <i>Namikawa T, Hanazaki K</i>
TOPIC HIGHLIGHT	2314	Non-classical phenotypes of autoimmune hepatitis and advances in diagnosis and treatment <i>Czaja AJ, Bayraktar Y</i>
ORIGINAL ARTICLES	2329	Signal transduction mechanism of TRB3 in rats with non-alcoholic fatty liver disease <i>Wang YG, Shi M, Wang T, Shi T, Wei J, Wang N, Chen XM</i>
BRIEF ARTICLES	2336	Non-steroidal anti-inflammatory drugs and statins in relation to colorectal cancer risk <i>Shadman M, Newcomb PA, Hampton JM, Wernli KJ, Trentham-Dietz A</i>
	2340	Study of the patency of different peritoneal drains used prophylactically in bariatric surgery <i>Salgado Júnior W, Macedo Neto MM, dos Santos JS, Sakarankutty AK, Ceneviva R, de Castro e Silva Jr O</i>
	2345	Celecoxib enhances the detoxification of diethylnitrosamine in rat liver cancer <i>Salcido-Neyoy ME, Sierra-Santoyo A, Beltrán-Ramírez O, Macías-Pérez JR, Villa-Treviño S</i>
	2351	Efficacy of the revised Vienna Classification for diagnosing colorectal epithelial neoplasias <i>Tominaga K, Fujinuma S, Endo T, Saida Y, Takahashi K, Maetani I</i>
	2357	Sclerosing cholangitis associated with autoimmune pancreatitis differs from primary sclerosing cholangitis <i>Kamisawa T, Takuma K, Anjiki H, Egawa N, Kurata M, Honda G, Tsuruta K</i>
	2361	Endoscopic ultrasonography does not differentiate neoplastic from non-neoplastic small gallbladder polyps <i>Cheon YK, Cho WY, Lee TH, Cho YD, Moon JH, Lee JS, Shim CS</i>
	2367	Mucin gene expression in bile of patients with and without gallstone disease, collected by endoscopic retrograde cholangiography <i>Vilkin A, Geller A, Levi Z, Niv Y</i>
2372	Determination of correlation of Adjusted Blood Requirement Index with outcome in patients presenting with acute variceal bleeding <i>Akhtar N, Zuberi BF, Hasan SR, Kumar R, Afsar S</i>	

- 2376** Local anesthesia with ropivacaine for patients undergoing laparoscopic cholecystectomy
Liu YY, Yeh CN, Lee HL, Wang SY, Tsai CY, Lin CC, Chao TC, Yeh TS, Jan YY
- 2381** Detection and evaluation of antibodies against neutrophil-activating protein of *Helicobacter pylori* in patients with gastric cancer
Long M, Luo J, Li Y, Zeng FY, Li M
- 2389** Association between Bmi1 and clinicopathological status of esophageal squamous cell carcinoma
He XT, Cao XF, Ji L, Zhu B, Lv J, Wang DD, Lu PH, Cui HG
- 2395** Polymorphisms of alcohol dehydrogenase-2 and aldehyde dehydrogenase-2 and esophageal cancer risk in Southeast Chinese males
Ding JH, Li SP, Cao HX, Wu JZ, Gao CM, Su P, Liu YT, Zhou JN, Chang J, Yao GH
- 2401** Diagnostic effect of capsule endoscopy in 31 cases of subacute small bowel obstruction
Yang XY, Chen CX, Zhang BL, Yang LP, Su HJ, Teng LS, Li YM
- 2406** Effect of two-channel gastric electrical stimulation with trains of pulses on gastric motility
Yang B, Hou XH, Song GQ, Liu JS, Chen JDZ

CASE REPORT

- 2412** Adult hereditary fructose intolerance
Yasawy MI, Folsch UR, Schmidt WE, Schwend M
- 2414** Drug-induced liver injury due to "natural products" used for weight loss: A case report
Tarantino G, Pezzullo MG, Dario di Minno MN, Milone F, Pezzullo LS, Milone M, Capone D
- 2418** Primary hepatic carcinoid: A case report and literature review
Fenoglio LM, Severini S, Ferrigno D, Gollè G, Serraino C, Bracco C, Castagna E, Brignone C, Pomero F, Migliore E, David E, Salizzoni M
- 2423** Biliary drainage of the common bile duct with an enteral metal stent
Dek IM, van den Elzen BDJ, Fockens P, Rauws EAJ
- 2425** Solitary extramedullary plasmacytoma in retroperitoneum: A case report and review of the literature
Hong W, Yu XM, Jiang MQ, Chen B, Wang XB, Yang LT, Zhang YP

ACKNOWLEDGMENTS

- 2428** Acknowledgments to reviewers of *World Journal of Gastroenterology*

APPENDIX

- 2429** Meetings
- 2430** Instructions to authors

FLYLEAF

- I-VII Editorial Board

INSIDE BACK COVER

Online Submissions

INSIDE FRONT COVER

Online Submissions

INTRODUCTION

World Journal of Gastroenterology is an international, open-access, peer-reviewed, and multi-disciplinary weekly journal that serves gastroenterologists and hepatologists. The biggest advantage of the open access model is that it provides free, full-text articles in PDF and other formats for experts and the public without registration, which eliminates the obstacle that traditional journals possess and usually delays the speed of the propagation and communication of scientific research results. The open access model has been proven to be a true approach that may achieve the ultimate goal of the journals, i.e. the maximization of the values of the readers, the authors and the society.

Maximization of the value of the readers can be comprehended in two ways. First, the journal publishes articles that can be directly read or downloaded free of charge at any time, which attracts more readers. Second, the readers can apply the knowledge in clinical practice without delay after reading and understanding the information in their fields. In addition, the readers are encouraged to propose new ideas based on those of the authors, or to provide viewpoints that are different from those of the authors. Such discussions or debates among different schools of thought will definitely boost advancements and developments in the fields. Maximization of the value of the authors refers to the fact that these journals provide a platform that promotes the speed of propagation and communication to a maximum extent. This is also what the authors really need. Maximization of the value of the society refers to the maximal extent of the social influences and impacts produced by the high quality original articles published in the journal. This is also the main purpose of many journals around the world.

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiao-Fang Liu*
Responsible Electronic Editor: *Yin-Ping Lin*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Lai-Fu Li*
Proofing Editorial Office Director: *Jian-Xia Cheng*

<p>NAME OF JOURNAL <i>World Journal of Gastroenterology</i></p> <p>RESPONSIBLE INSTITUTION Department of Science and Technology of Shanxi Province</p> <p>SPONSOR Taiyuan Research and Treatment Center for Digestive Diseases, 77 Shuangta Xijie, Taiyuan 030001, Shanxi Province, China</p> <p>EDITING Editorial Board of <i>World Journal of Gastroenterology</i>, Room 903, Building D, Ocean International Center, No.62 Dongsihuan Zhonglu, Chaoyang District, Beijing 100025, China Telephone: +86-10-59080039 Fax: +86-10-85381893 E-mail: wjg@wjgnet.com http://www.wjgnet.com</p> <p>PUBLISHING The WJG Press and Beijing Baishideng BioMed Scientific Co., Ltd., Room 903, Building D, Ocean International Center, No.62 Dongsihuan Zhonglu, Chaoyang District, Beijing 100025, China Telephone: +86-10-59080039 Fax: +86-10-85381893 E-mail: wjg@wjgnet.com http://www.wjgnet.com</p> <p>PRINTING Beijing Kexin Printing House</p> <p>OVERSEAS DISTRIBUTOR Beijing Bureau for Distribution of Newspapers and Journals (Code No. 82-261) China International Book Trading Corporation PO Box 399, Beijing, China (Code No. M4481)</p> <p>PUBLICATION DATE May 21, 2009</p> <p>EDITOR-IN-CHIEF Lian-Sheng Ma, <i>Beijing</i></p>	<p>SUBSCRIPTION RMB 50 Yuan for each issue, RMB 2400 Yuan for one year</p> <p>CSSN ISSN 1007-9327 CN 14-1219/R</p> <p>HONORARY EDITORS-IN-CHIEF Montgomery Bissell, <i>San Francisco</i> James L Boyer, <i>New Haven</i> Chao-Long Chen, <i>Kaohsiung</i> Ke-Ji Chen, <i>Beijing</i> Li-Fang Chou, <i>Taipei</i> Jacques V Dam, <i>Stanford</i> Martin H Floch, <i>New Haven</i> Guadalupe Garcia-Tsao, <i>New Haven</i> Zhi-Qiang Huang, <i>Beijing</i> Shinn-Jang Hwang, <i>Taipei</i> Ira M Jacobson, <i>New York</i> Derek Jewell, <i>Oxford</i> Emmet B Keeffe, <i>Palo Alto</i> Min-Liang Kuo, <i>Taipei</i> Nicholas F LaRusso, <i>Rochester</i> Jie-Shou Li, <i>Nanjing</i> Geng-Tao Liu, <i>Beijing</i> Lein-Ray Mo, <i>Tainan</i> Bo-Rong Pan, <i>Xi'an</i> Fa-Zu Qiu, <i>Wuhan</i> Eamonn M Quigley, <i>Cork</i> David S Rampton, <i>London</i> Rafiq A Sheikh, <i>Sacramento</i> Rudi Schmid, <i>Kentfield</i>¹⁾ Nicholas J Talley, <i>Rochester</i> Sun-Lung Tsai, <i>Yong-Kang City</i> Guido NJ Tytgat, <i>Amsterdam</i> Hsiu-Po Wang, <i>Taipei</i> Jaw-Ching Wu, <i>Taipei</i> Meng-Chao Wu, <i>Shanghai</i> Ming-Shiang Wu, <i>Taipei</i> Jia-Yu Xu, <i>Shanghai</i> Ta-Sen Yeh, <i>Taoyuan</i> Ming-Lung Yu, <i>Kaohsiung</i></p> <p>STRATEGY ASSOCIATE EDITORS-IN-CHIEF Peter Draganov, <i>Florida</i> Ronnie Fass, <i>Tucson</i> Hugh J Freeman, <i>Vancouver</i> John P Geibel, <i>New Haven</i> Maria C Gutiérrez-Ruiz, <i>México</i></p>	<p>Kazuhiro Hanazaki, <i>Kochi</i> Akio Inui, <i>Kagoshima</i> Kalpesh Jani, <i>Vadodara</i> Sanaa M Kamal, <i>Cairo</i> Ioannis E Koutroubakis, <i>Heraklion</i> Jose JG Marin, <i>Salamanca</i> Javier S Martin, <i>Punta del Este</i> Natalia A Osna, <i>Omaha</i> Jose Sahel, <i>Marseille</i> Ned Snyder, <i>Galveston</i> Nathan Subramaniam, <i>Brisbane</i> Wei Tang, <i>Tokyo</i> Alan BR Thomson, <i>Edmonton</i> Paul Joseph Thuluvath, <i>Baltimore</i> James F Trotter, <i>Denver</i> Shingo Tsuji, <i>Osaka</i> Harry HX Xia, <i>Hanover</i> Yoshio Yamaoka, <i>Houston</i> Jesus K Yamamoto-Furusho, <i>México</i></p> <p>ASSOCIATE EDITORS-IN-CHIEF Gianfranco D Alpini, <i>Temple</i> Bruno Annibale, <i>Roma</i> Roger William Chapman, <i>Oxford</i> Chi-Hin Cho, <i>Hong Kong</i> Alexander L Gerbes, <i>Munich</i> Shou-Dong Lee, <i>Taipei</i> Walter Edwin Longo, <i>New Haven</i> You-Yong Lu, <i>Beijing</i> Masao Omata, <i>Tokyo</i></p> <p>EDITORIAL OFFICE Director: Jian-Xia Cheng, <i>Beijing</i> Deputy Director: Jian-Zhong Zhang, <i>Beijing</i></p> <p>LANGUAGE EDITORS Director: Jing-Yun Ma, <i>Beijing</i> Deputy Director: Xian-Lin Wang, <i>Beijing</i></p> <p>MEMBERS Gianfranco D Alpini, <i>Temple</i> BS Anand, <i>Houston</i> Manoj Kumar, <i>Nepal</i> Patricia F Lalor, <i>Birmingham</i> Ming Li, <i>New Orleans</i> Margaret Lutze, <i>Chicago</i> Sabine Mihm, <i>Göttingen</i> Francesco Negro, <i>Genève</i> Bernardino Rampone, <i>Siena</i> Richard A Rippe, <i>Chapel Hill</i> Stephen E Roberts, <i>Swansea</i></p>	<p>COPY EDITORS Gianfranco D Alpini, <i>Temple</i> Sujit Kumar Bhattacharya, <i>Kolkata</i> Filip Braet, <i>Sydney</i> Kirsteen N Browning, <i>Baton Rouge</i> Radha K Dhiman, <i>Chandigarh</i> John Frank Di Mari, <i>Texas</i> Shannon S Glaser, <i>Temple</i> Eberhard Hildt, <i>Berlin</i> Patricia F Lalor, <i>Birmingham</i> Ming Li, <i>New Orleans</i> Margaret Lutze, <i>Chicago</i> MI Torrs, <i>Jaén</i> Sri Prakash Misra, <i>Allahabad</i> Bernardino Rampone, <i>Rome</i> Giovanni Monteleone, <i>Rome</i> Valerio Nobili, <i>Rome</i> Osman Cavit Ozdogan, <i>Istanbul</i> Francesco Perri, <i>San Giovanni Rotondo</i> Thierry Piche, <i>Nice</i> Bernardino Rampone, <i>Siena</i> Richard A Rippe, <i>Chapel Hill</i> Ross C Smith, <i>Sydney</i> Daniel Lindsay Worthley, <i>Bedford</i> George Y Wu, <i>Farmington</i> Jian Wu, <i>Sacramento</i></p> <p>COPYRIGHT © 2009 Published by The WJG Press and Baishideng. All rights reserved; no part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior permission of WJG. Authors are required to grant WJG an exclusive licence to publish.</p> <p>SPECIAL STATEMENT All articles published in this journal represent the viewpoints of the authors except where indicated otherwise.</p> <p>INSTRUCTIONS TO AUTHORS Full instructions are available online at http://www.wjgnet.com/wjg/help/instructions.jsp. If you do not have web access please contact the editorial office.</p> <p>ONLINE SUBMISSION http://wjg.wjgnet.com</p>
--	---	---	--

Determination of correlation of Adjusted Blood Requirement Index with outcome in patients presenting with acute variceal bleeding

Naheed Akhtar, Bader Faiyaz Zuberi, Syed Riazul Hasan, Raj Kumar, Salahuddin Afsar

Naheed Akhtar, Bader Faiyaz Zuberi, Syed Riazul Hasan, Raj Kumar, Salahuddin Afsar, Department of Medicine, Dow University of Health Sciences, Karachi 74000, Pakistan
Author contributions: Zuberi BF designed the study and carried out the data/statistical analysis; Akhtar N, Hasan SR and Kumar R contributed equally to data collection and manuscript writing; Afsar S performed the final editing and review of the manuscript.

Correspondence to: Dr. Bader Faiyaz Zuberi, Department of Medicine, Dow University of Health Sciences, Karachi 74000, Pakistan. bader@zuberi.net

Telephone: +92-300-8234883 Fax: +92-21-9216027

Received: March 7, 2009 Revised: April 23, 2009

Accepted: April 30, 2009

Published online: May 21, 2009

© 2009 The WJG Press and Baishideng. All rights reserved.

Key words: Adjusted Blood Requirement Index; Cirrhosis; Mortality; Portal hypertension; Variceal hemorrhage

Peer reviewer: Abdellah Essaid, Professor, Hospital Ibn Sina, Rabat 10100, Morocco

Akhtar N, Zuberi BF, Hasan SR, Kumar R, Afsar S. Determination of correlation of Adjusted Blood Requirement Index with outcome in patients presenting with acute variceal bleeding. *World J Gastroenterol* 2009; 15(19): 2372-2375 Available from: URL: <http://www.wjgnet.com/1007-9327/15/2372.asp> DOI: <http://dx.doi.org/10.3748/wjg.15.2372>

Abstract

AIM: To determine the correlation of Adjusted Blood Requirement Index (ABRI) with the 7th day outcome in patients presenting with acute variceal bleeding.

METHODS: All patients presenting with acute variceal hemorrhage (AVH) were included. Patients with previous band ligation, sclerotherapy, gastrointestinal or hepatic malignancies were excluded. Patients were managed as per standard protocol for AVH with terlipressin and band ligation. ABRI scores were calculated using the formula outcome of alive or expired up to the 7th day after treatment. The correlation between ABRI and mortality was estimated and a receiver operative characteristic (ROC) curve was plotted.

RESULTS: A total of 113 patients (76 male; 37 female) were included. On assessment, 18 were in Child's Pugh Class A, 82 in Class B and 13 were in Class C. The median number of blood units transfused \pm inter-quartile range was 3.0 ± 2.0 . The median \pm inter-quartile range for ABRI was 1.3 ± 1.1 . The ROC curve of ABRI for expiry showed a significantly large area of 0.848 ($P < 0.0001$; 95% CI: 0.75-0.95). A significant correlation of log transformation of ABRI with an outcome of mortality was present ($P < 0.0001$).

CONCLUSION: ABRI correlates strongly with mortality.

INTRODUCTION

Chronic liver diseases and cirrhosis are now being recognized as an important cause of morbidity and mortality worldwide. Acute variceal hemorrhage (AVH) secondary to cirrhosis is to date the most important cause of mortality in cirrhosis^[1]. In Pakistan, hepatitis B and C are the most important causes of cirrhosis^[2]. The frequency of varices is very high in cirrhotic patients, nearly 40% of patients with compensated cirrhosis and 60% with decompensated cirrhosis have varices^[3]. Due to recent advancements, mortality from AVH has been reduced to 20% from the first variceal bleed^[4]. Bleeding from AVH carries a high risk of mortality during the first 5 d, with a gradual decline in risk over the next 4-6 wk^[5]. The prediction and evaluation of adequate hemostasis by non-endoscopic methods are desired by treating physicians. Many criteria and definitions to evaluate failure to control and prevent variceal bleeding were developed in the Baveno Consensus Workshops I - III but failed in clinical application due to cumbersome procedures and calculations^[6-10]. Further developments in this subject identified an independent factor the "Adjusted Blood Requirement Index (ABRI)" in the Baveno Workshop IV^[11]. ABRI was developed to determine adequate control or failure to control variceal hemorrhage. An ABRI value of ≥ 0.75 at any point time was defined as a failure to control variceal bleeding^[11].

Failure of AVB control leads to increased mortality. Thus ABRI could be used to assess the risk of mortality. A correlation between ABRI and mortality has not been evaluated in a prospectively designed study. We have reported its correlation with outcome in a retrospective analysis previously^[12]. As there are no reports of a prospective evaluation of ABRI and its relation to mortality, there is a need to assess this correlation in our settings.

This study was designed to evaluate the correlation between ABRI and outcome at the 7th day after hospital admission as improved or expired in acute variceal bleeding.

MATERIALS AND METHODS

All cirrhotic patients who presented with AVB were included. Informed consent was obtained from all patients. Patients with a history of previous band ligation or sclerotherapy, hepatocellular carcinoma and the presence of peptic ulcer or gastrointestinal (GI) malignancy on endoscopy were excluded. Patients were managed as per standard protocol of acute variceal bleeding^[10]. Blood samples were taken for Complete Blood Counts, Prothrombin Time, Liver Function Tests and albumin before the start of therapy. Child's Pugh Class assessment was carried out. All patients were given terlipressin 2 mg *iv* initial dose and followed by 1 mg/6 h for 3 d. The number of blood units transfused was noted and endoscopic variceal band ligation (EVBL) was performed within 24 h of admission. Study endpoint was patient outcome (alive or expired at the 7th day after admission). The ABRI value was calculated using the following formula^[11]: $ABRI = \text{blood units transfused} / [(\text{final hematocrit} - \text{initial hematocrit}) + 0.01]$. Child's Pugh score was calculated using the formula^[13] shown in Table 1.

Sample size

Sample size was estimated using the following parameters: Level of Significance (α) = 5%; Power of test ($1 - \beta$) = 80%; Test value of population proportion (P_0) = 20% (0.2); Anticipated value of population proportion (P_a) = 30% (0.3); Sample size (n) = 109.

Statistical analysis

mean \pm SD was calculated for age. Median and inter-quartile range were calculated for the number of blood units transfused and ABRI. Frequencies of gender, Child's Pugh Class and outcome were calculated. ABRI values ≥ 0.75 were recoded into a new variable as uncontrolled while ABRI values < 0.75 were recoded as controlled and their frequency estimated. χ^2 test was performed for outcome with ABRI control status and Child's Pugh Class was carried out with continuity correction and likelihood ratio applied where indicated. A receiver operative characteristic (ROC) curve of ABRI was plotted for expiry. Log transformation of variable ABRI was carried out as it was not normally distributed and then used for Pearson's Bivariate correlation with outcome. The significance level was set

Table 1 Child's Pugh score was calculated (using formula)

Parameter	Numerical score		
	1	2	3
Ascites	None	Slight	Moderate to severe
Encephalopathy	None	Slight to moderate	Moderate to severe
Bilirubin (mg/dL)	< 2.0	2-3	> 3.0
Albumin (g/dL)	> 3.5	2.8-3.5	< 2.8
Prothrombin time (prolonged in seconds)	1-3 s	4-6 s	> 6.0

Child's Pugh Class A = 5-6 points; Child's Pugh Class B = 7-9 points; Child's Pugh Class C = 10-15 points.

Table 2 Cross tabulation of ABRI groups with outcome

ABRI groups		Outcome		Total
		Alive	Expired	
ABRI groups	Controlled	27	0	27
	Uncontrolled	67	19	86
Total		94	19	113

at $P \leq 0.05$. SPSS version 17.0 was used for statistical analysis.

RESULTS

A total of 113 patients fulfilling the inclusion/exclusion criteria were inducted. These included 76 (67.3%) male (44.3 ± 11.8 years) and 37 (32.7%) female (44.1 ± 9.4 years). Terlipressin was given to 111 patients (98.2%) immediately on admission. EVBL was performed in 105 (92.9%) patients. The assessment on admission showed that 18 (15.9%) were in Child's Pugh Class A; 82 (72.6%) in Child's Pugh Class B and 13 (11.5%) were in Child's Pugh Class C. The median number of blood transfusions given was 3.0 pints and the inter-quartile range was 2.0. The median ABRI was 1.3 with an inter-quartile range of 1.1. The number of patients with $ABRI \geq 0.75$ was 86 (76.1%) showing a failure to control variceal bleeding according to ABRI criteria. Outcome at the 7th day after admission showed that 94 (83.2%) patients were alive while 19 (16.8%) patients had expired during this period. Cross tabulation of outcome (alive and expired) with ABRI status [controlled (< 0.75) and uncontrolled (≥ 0.75)] showed that no patients had expired in the ABRI controlled group (Table 2). χ^2 test with continuity correction gave a significance value of $P = 0.017$. A similar cross tabulation with Child's Pugh Class showed that the highest percentage of patients expired in Child's Pugh Class C while no patients with Child's Class A expired (Table 3). χ^2 test with the Likelihood Ratio gave significant differences in the frequencies of expiry with Child's Pugh Class ($P < 0.0001$). A ROC curve was plotted using expiry as a state variable (Figure 1). The area under the curve was significantly large at 0.848 ($P < 0.0001$; 95% CI: 0.75-0.95). The sensitivity and specificity of the ABRI cutoff value of 0.75 in our study was 100% and 73.4%, respectively. The correlation of

Table 3 Cross tabulation of Child's Pugh Class with outcome

			Outcome		Total
			Alive	Expired	
Child's Pugh Class	Class A	Count	18	0	18
		% within Child's Pugh Class	100.0%	0.0%	100.0%
	Class B	Count	71	11	82
		% within Child's Pugh Class	86.6%	13.4%	100.0%
	Class C	Count	5	8	13
		% within Child's Pugh Class	38.5%	61.5%	100.0%
Total		Count	94	19	113
		% within Child's Pugh Class	83.2%	16.8%	100.0%




Figure 1 ROC curve of ABRI against expiry.

ABRI with outcome was analyzed by Pearson's Bivariate Correlation test. As the data of ABRI was skewed and not normally distributed its Log_{10} transformation was used. The results showed a significant correlation between ABRI and expiry with $P < 0.0001$.

DISCUSSION

This study showed a significant correlation between ABRI and the 7th day outcome in patients with AVH. This is a very significant finding as it is important to predict the outcome at an initial stage of management and ABRI suggests whether the variceal hemorrhage has been arrested at any point during management. Our study also established its correlation with mortality. Earlier validation studies of ABRI were retrospective^[11,12]. The current study is prospective and designed more specifically to assess the correlation of ABRI with mortality which has not been previously studied. The correlation of higher ABRI scores with mortality was significant and this simple to use parameter should be used to assess failure to control bleeding and risk of mortality. The number of units of blood transfused and hematocrit levels, if used alone, are not good criteria to assess variceal bleeding control. We also used pharmacological and endoscopic interventions and the combined effect of these interventions was reflected in the outcome which was also observed in other reports from this region^[14]. About 70% of patients rebleeding within 2 years, thus managing the index bleed properly and obliteration of varices can decrease rebleeding^[15,16].

Many scoring systems have been derived to predict the outcome of upper GI hemorrhage. The Rockall score is one such scoring system for predicting rebleeding and mortality which also showed good correlation^[17,18]. Limitations of the Rockall score are that it is rather difficult to use with the requirement of more parameters as compared to ABRI and it is not variceal bleeding specific, but designed for both variceal and non-variceal bleeding^[18,19]. Another popular scoring system, the Child's Pugh score predicts all cause morbidity and mortality in cirrhotic patients but is not specific for variceal hemorrhage^[20].

In conclusion, among the many predictive scoring systems in cirrhotic patients, ABRI is specific for variceal hemorrhage and correlates strongly with mortality and is a good indicator of the failure of variceal hemorrhage control.

COMMENTS

Background

In a developing country like Pakistan, hepatitis B and C are the most important causes of cirrhosis. The frequency of varices is very high in cirrhotic patients, nearly 40% of patients with compensated cirrhosis and 60% of patients with decompensated cirrhosis have varices. Mortality from variceal bleeding is still high at 20%. The prediction of mortality risk is difficult and available scores are difficult to calculate and thus do not enjoy wide acceptability and application.

Research frontiers

Adjusted Blood Requirement Index (ABRI) is a score which is used to determine the failure to control variceal bleeding. In this study it was correlated with the outcome of mortality.

Innovations and breakthroughs

Many scoring systems are in use to predict the outcome of upper gastrointestinal hemorrhage. The Rockall score is one such scoring system for predicting rebleeding and mortality but it is rather difficult to use. Another scoring system, the Child's Pugh score predicts all cause morbidity and mortality in cirrhotic patients but is not specific for variceal hemorrhage. The ABRI is a variceal hemorrhage-specific score and is easy to use.

Applications

ABRI: A validated tool to determine variceal bleeding control also correlates well with mortality in such patients.

Peer review

Many scoring systems have been described to predict the prognosis of upper gastrointestinal hemorrhage like Rockall and Child's Pugh but these have limitations. In practice, it is useful to predict the outcome at admission of patients with acute variceal hemorrhage. This study showed a significant correlation between ABRI and expiry. The methodology is correct. This work deserves to be published to stimulate other teams over the world to perform the same study with a large number of patients.

REFERENCES

- 1 **Burroughs AK**, Triantos CK, O'Beirne J, Patch D. Predictors of early rebleeding and mortality after acute variceal hemorrhage in patients with cirrhosis. *Nat Clin Pract Gastroenterol Hepatol* 2009; **6**: 72-73
- 2 **Mashud I**, Khan H, Khattak AM. Relative frequency of hepatitis B and C viruses in patients with hepatic cirrhosis at DHQ Teaching Hospital D. I. Khan. *J Ayub Med Coll Abbottabad* 2004; **16**: 32-34
- 3 **Schepis F**, Cammà C, Niceforo D, Magnano A, Pallio S, Cinquegrani M, D'amico G, Pasta L, Craxì A, Saitta A, Raimondo G. Which patients with cirrhosis should undergo endoscopic screening for esophageal varices detection? *Hepatology* 2001; **33**: 333-338
- 4 **Turnes J**, Garcia-Pagan JC, Abraldes JG, Hernandez-Guerra M, Dell'Era A, Bosch J. Pharmacological reduction of portal pressure and long-term risk of first variceal bleeding in patients with cirrhosis. *Am J Gastroenterol* 2006; **101**: 506-512
- 5 **Burroughs AK**, Mezzanotte G, Phillips A, McCormick PA, McIntyre N. Cirrhotics with variceal hemorrhage: the importance of the time interval between admission and the start of analysis for survival and rebleeding rates. *Hepatology* 1989; **9**: 801-807
- 6 **North Italian Endoscopic Club for the Study and Treatment of Esophageal Varices**. Prediction of the first variceal hemorrhage in patients with cirrhosis of the liver and esophageal varices. A prospective multicenter study. *N Engl J Med* 1988; **319**: 983-989
- 7 **de Franchis R**. Developing consensus in portal hypertension. *J Hepatol* 1996; **25**: 390-394
- 8 **de Franchis R**. Updating consensus in portal hypertension: report of the Baveno III Consensus Workshop on definitions, methodology and therapeutic strategies in portal hypertension. *J Hepatol* 2000; **33**: 846-852
- 9 **de Franchis R**. Evaluation and follow-up of patients with cirrhosis and oesophageal varices. *J Hepatol* 2003; **38**: 361-363
- 10 **de Franchis R**. Evolving consensus in portal hypertension. Report of the Baveno IV consensus workshop on methodology of diagnosis and therapy in portal hypertension. *J Hepatol* 2005; **43**: 167-176
- 11 **Duvnjak M**, Barsić N, Tomasić V, Jukić LV, Lerotić I, Pavić T. Adjusted blood requirement index as indicator of failure to control acute variceal bleeding. *Croat Med J* 2006; **47**: 398-403
- 12 **Zuberi BF**, Riaz MF, Sultan BA, Gobindram P, Farooq A, Qadeer R, Memon AR, Afsar S. Correlation of adjusted blood requirement index with treatment intervention and outcome in patients presenting with acute variceal bleeding. *J Dow Uni Health Sci* 2007; **1**: 65-68
- 13 **D'Amico G**, Garcia-Tsao G, Pagliaro L. Natural history and prognostic indicators of survival in cirrhosis: a systematic review of 118 studies. *J Hepatol* 2006; **44**: 217-231
- 14 **Varghese J**, Cherian JV, Solomon R, Jayanthi V. Predictors of variceal bleed among patients with liver cirrhosis in the era of sclerotherapy. *Singapore Med J* 2008; **49**: 239-242
- 15 **Bureau C**, Vinel JP. Management of failures of first line treatments. *Dig Liver Dis* 2008; **40**: 343-347
- 16 **Bosch J**, Abraldes JG, Berzigotti A, Garcia-Pagan JC. Portal hypertension and gastrointestinal bleeding. *Semin Liver Dis* 2008; **28**: 3-25
- 17 **Sarwar S**, Dilshad A, Khan AA, Alam A, Butt AK, Tariq S, Ahmad I. Predictive value of Rockall score for rebleeding and mortality in patients with variceal bleeding. *J Coll Physicians Surg Pak* 2007; **17**: 253-256
- 18 **Sanders DS**, Carter MJ, Goodchap RJ, Cross SS, Gleeson DC, Lobo AJ. Prospective validation of the Rockall risk scoring system for upper GI hemorrhage in subgroups of patients with varices and peptic ulcers. *Am J Gastroenterol* 2002; **97**: 630-635
- 19 **Tham TC**, James C, Kelly M. Predicting outcome of acute non-variceal upper gastrointestinal haemorrhage without endoscopy using the clinical Rockall Score. *Postgrad Med J* 2006; **82**: 757-759
- 20 **Sarwar S**, Khan AA, Tarique S. Comparison of MELD, Child Pugh score and Rockall score for predicting rebleeding and in-hospital mortality in patients of variceal bleeding. *J Coll Physicians Surg Pak* 2008; **18**: 524-525

S- Editor Tian L L- Editor Webster JR E- Editor Zheng XM